




SCIENCE SPOTLIGHT™

THE 3CL_{pro} INHIBITOR ALG-097111 POTENTLY INHIBITS SARS-CoV-2 REPLICATION IN HAMSTERS


Koen Vandyck

*On behalf of the Aligos/CD3/Rega collaboration team
Aligos Belgium BV, Leuven, Belgium*


Disclosure: Koen Vandyck is an employee of Aligos Belgium BV

SARS-CoV-2 3CL_{pro} Represents an Attractive Target

- The main protease of SARS-CoV-2 3CL_{pro} is conserved among coronaviruses (including SARS-CoV-1, 2 and MERS-CoV)
- 3CL_{pro} cleaves at multiple sites on the large polyprotein 1ab
- No human homolog of 3CL_{pro}
- Collaboration on Coronavirus Protease Inhibitors between Aligos and CD3/Rega at KU Leuven



Other Reported 3CL_{pro} Inhibitors also Inhibit Cathepsin L



	Cpd 11a	6j	6e	Cpd 13b	Cpd 11r	A9	PF-00835231
Cathepsin L IC ₅₀ (nM)	0.21 (n=2)	<0.5 (n=2)	<0.5 (n=2)	291 (n=2)	<0.5 (n=2)	<0.5 (n=2)	146 (n=13)
SARS-CoV-2 3CL _{pro} IC ₅₀ (nM)	8 (n=28)	7 (n=2)	10 (n=3)	472 (n=2)	1154 (n=2)	4891 (n=2)	5 (n=5)

- Cathepsin L is involved in SARS-CoV-2 entry and inhibitors block replication in several cell lines
- The SARS-CoV-2 inhibitory activity of Cathepsin L inhibitors is reduced upon expression of TMPRSS2 and might not translate *in vivo*


Increasing the selectivity of SARS-CoV-2 3CL_{pro} inhibitors over Cathepsin L is considered important

ALG-097111 is a Selective SARS-CoV-2 3Cl_{pro} Inhibitor




Biochemical

SARS-CoV-2 3CL_{pro} (**IC₅₀ = 7 nM**)
human cathepsin L (**IC₅₀ > 10 μM**)




Cellular

SARS-CoV-2 (A549+ACE2): **EC₅₀ = 200 nM**
(similar potency confirmed for β-CoV OC43 (HeLa) and α-CoV 229E (Huh-7))



SARS-CoV-2 inhibition in human small airway epithelial cell cultures
>3 log₁₀ viral RNA reduction




ALG-097111, a selective SARS-CoV-2 3Cl_{pro} inhibitor, does not inhibit Cathepsin L, and shows cellular potency vs. alpha and beta coronaviruses

ALG-097111 Dosing in Female Syrian Hamster


- ALG-097111 is stable in human and dog liver microsomes ($t_{1/2}$ >60 min) and hepatocytes ($t_{1/2}$ >360 min)
- ALG-097111 is stable in hamster liver microsomes only in the presence of ritonavir ($t_{1/2}$ >60 min)
- Dose in hamsters: 200 mg/kg/dose BID SC co-dosed with ritonavir PO 50 mg/kg/dose BID
- Hamster plasma and lung C_{trough} : 5.5 and 3.0-fold above SARS-CoV-2 A549-ACE2 EC_{50}

ALG-097111 Plasma PK profile and lung concentration in female hamsters




SC administration of ALG-097111 when co-dosed with ritonavir, resulted in significant systemic exposure in female Syrian hamsters

ALG-097111 Efficacy in SARS-CoV-2 Hamster Model



Infectious viral loads in the lungs of SARS-CoV-2-infected hamsters at day 2 post-infection



ALG-097111, a selective SARS-CoV-2 3CL_{pro} inhibitor efficiently inhibits viral replication in the lungs of infected hamsters